

# Scaling Basic Toxicokinetic Parameters from Rat to Man

Kenneth Bachmann,<sup>1</sup> David Pardoe,<sup>2</sup> and Donald White<sup>1,2</sup>

<sup>1</sup>Department of Pharmacology and <sup>2</sup>Department of Mathematics, The University of Toledo, Toledo, OH 43606 USA

Scaling of the quantified dispositional parameters of xenobiotics from animals to man is of interest from the standpoint of toxicology (e.g., poisoning and risk assessment). Scaling is also important from the standpoint of therapeutics because it represents a strategy for predicting first-use-in-human doses in clinical trials of investigational new drugs. Current strategies for scaling either doses of xenobiotics or the dispositional parameters of xenobiotics from animals to man rely on models that take account principally of species differences in weight or body surface area. Interspecies scaling of dispositional parameters such as clearance or volume of distribution commonly involves the comparison of estimates of these parameters for a given xenobiotic among numerous species on the basis of weight with the resultant mathematical relationship used to predict the values of those parameters for that xenobiotic in a species weighing, on average, about 70 kg (i.e., a man). Our approach has been to ascertain whether a useful mathematical model could be developed for predicting the dispositional parameters of a xenobiotic, its half-life and volume of distribution, in humans based exclusively on estimates of those parameters in rats. Based on a data set of about 100 different xenobiotics, we found that values for half-life and volume of distribution of a xenobiotic in humans can be predicted from the estimates of those parameters in rats. **Key words:** half-life, humans, pharmacokinetics, rats, scaling, toxicokinetics, volume of distribution. *Environ Health Perspect* 104:400–407 (1996)

Predicting the dispositional kinetics of xenobiotics in humans from data acquired in animals is an important enterprise. Predicted pharmacokinetic parameters have application in drug development such as in the prediction of initial doses of investigational new drugs for human studies. Predicted toxicokinetic parameters have applications in toxicology such as in the prediction of the time course of exposure after acute or chronic challenge with toxic substances. Knowledge of the time course may be helpful in predicting both the aggressiveness of treatments and the duration of such treatments. Another application is in carcinogenic risk assessment.

Estimates of toxicokinetic parameters in humans for individual xenobiotics may be attained either through interspecies allometric scaling (1,2) or through physiologically based pharmacokinetic (PBPK) modeling (3), which ultimately must invoke an interspecies scale-up as well, at least for data elicited exclusively from an animal model (4). In allometric scaling, kinetic variables such as volume of distribution and half-life are scaled based on the assumption that they are proportional to a power of body weight, and xenobiotic clearance is additionally factored by the species' life span (5).

Interspecies scaling as it relates to dosimetry centers on adjustments based on species mass (i.e., body weight), though some prefer that adjustments from one species to another take account of body surface area rather than body weight. For example, the U.S. Environmental Protection Agency suggests that doses be scaled

from rats to humans on the basis of the ratio of human body weight to rat body weight (BW) raised to the 0.667 power, as follows (6):

$$\text{Dose}_{\text{human}} = \text{Dose}_{\text{rat}} (\text{BW}_{\text{human}} / \text{BW}_{\text{rat}})^{0.667} \quad (1)$$

Recent proposals suggest that the exponent used should be 0.75 (7) rather than 0.667, and this value appears to be justifiable on the basis that basal metabolic rate is a function of mass raised to the 0.75 power as shown (8):

$$\text{Basal metabolism} = a M^{0.75} \quad (2)$$

where  $a$  is a mass coefficient set at 70 kcal and  $M$  is mass in kilograms. Related parameters that scale to  $\text{BW}^{0.75}$  include alveolar ventilation, cardiac output, renal clearance, and oxygen consumption (9). Most often, the allometric scaling of kinetic variables is undertaken on a case-by-case basis (i.e., xenobiotic by xenobiotic), with kinetic parameters for individual xenobiotics evaluated in each of multiple species (10).

We were interested in determining whether reasonable predictions of two kinetic parameters of xenobiotics, volume of distribution and half-life, could be made for humans if one had knowledge of those parameters from rat data only. Our working hypothesis was that a useful predictive mathematical relationship was likely to exist if one regressed known human toxicokinetic variable estimates (volume of distribution and half-life) against known data for the same variables in rats, and did

so across a large series of xenobiotics. Such a relationship would establish a single model from which volume of distribution and half-life for any xenobiotic could be predicted for humans and, further, could be predicted entirely from the estimates of those variables in rats.

## Methods

Average values for rats and humans for volumes of distribution of 100 xenobiotics and for half-lives of 103 xenobiotics were obtained from the literature. Most of the xenobiotics were "drugs" insofar as there is a substantially larger literature for human pharmacokinetic parameters than toxicokinetic parameters. A table denoting each xenobiotic, the average toxicokinetic or pharmacokinetic parameters for both rats and humans and the source of the data are given in the appendix.

We modeled half-life and volume of distribution data using the statistical computation program, S-Plus (Mathsoft Inc., Seattle, Washington), which provided confidence limits for model predictions. The program ran on a SPARCserver 10 computer. Our aim was to find a simple yet effective model describing the human data as a function of that of the rat. Further, it was important to identify confidence limits on predictions that may arise from such a model. Noting how the variability in each data set increases linearly with the values for the rat for each parameter we decided to explore logarithmic models because these models are inherently multiplicative in the error terms, as will be seen below.

## Results

After confirming that quadratic terms were not statistically significant, and upon noting that the logarithmic transformations yielded a standard deviation for human values that was constant across rat values (see Figs. 1 and 2), a linear model that is logarithmic in both variables was selected for both data sets:

$$\log(P_{i_{\text{human}}}) = \alpha + [b \times \log(P_{i_{\text{rat}}})] + e_i, \quad i = 1, 2, \dots, n \quad (3)$$

Address correspondence to K. Bachmann, Department of Pharmacology, The University of Toledo College of Pharmacy, 2801 W. Bancroft Street, Toledo, OH 43606 USA.

This work was supported in part by a grant from the Gustavus Pfeiffer Foundation.

Received 27 September 1995; accepted 18 December 1995.


where  $P$  denotes the kinetic parameter (half-life or volume of distribution) and the errors,  $e_i$ , are assumed to be normally distributed with a constant standard deviation,  $\sigma$ . This model can be rewritten in more familiar terms as:

$$P_{i_{\text{human}}} = a_i \times P_{i_{\text{rat}}}^b \quad (4)$$


where  $a_i$  depends on both the parameter  $\alpha$  and the individual errors,  $e_i$ . Thus, the variability of the actual human parameter value is proportional to the rat value raised to the power  $b$ .

Results of the regression analysis from S-Plus are given in Table 1. Included are the parameter estimates along with their standard errors for each data set. All the parameters are significant with  $p$ -values based on the  $t$ -distribution  $<0.0001$ . The  $R^2$  values are 0.737 and 0.754 for half-life and volume of distribution analyses, respectively, demonstrating a reasonably good fit in both cases. The estimated regression curves, or prediction lines, for half-life and volume of distribution are shown in Figures 1 and 2, respectively. The exponents ( $b$ ) are 0.83 for half-life and 0.91 for volume of distribution, respectively. Also given in these figures are the 80%, 90%, and 95% prediction intervals for a new human value corresponding to the given rat value. These curves take into account the possible errors in the prediction line itself (in the estimated  $\alpha$  and  $b$ ) and the error inherent to a new observation (as quantified by the regression estimate of  $\sigma$ ).

The validity of the regression equations was confirmed as follows: for the half-life data, the regression equation along with the prediction intervals was computed using a subset of the data originally obtained (i.e., all but 18 of the data points ultimately used). The additional 18 data points (these are marked with footnote *a* in the data listings in the appendix) obtained later were compared to the regression results with these points excluded. These points followed basically the same pattern as the larger data set. Furthermore, two of these points fell outside the 80% prediction intervals (about three or four would have been expected to do so); none of these points fell outside the 90% prediction intervals (one or two would have been predicted to do so); and none fell outside the 95% prediction intervals (one or fewer would be expected). As the regression relationship thus proved to be useful in predicting human half-lives from rat half-lives, we concluded that the regression results obtained were valid. Note that the regression results in the tables and figures employ



**Figure 1.** Human half-lives for 103 xenobiotics plotted against rat half-lives. Half-lives are given in hours. Both axes are logarithmically scaled.



**Figure 2.** Human volumes of distribution for 100 xenobiotics plotted against rat volumes of distribution. Volumes of distribution are given in liters/kilogram. Both axes are logarithmically scaled.

**Table 1.** Regression analysis for human versus rat parameters

	Estimate	SE	<i>t</i>	<i>p</i>	Residual SE	Multiple <i>R</i> <sup>2</sup>
Half-life						
Intercept ( $\alpha$ )	1.58	0.083	19.09	0.0000		
Slope ( $b$ )	0.83	0.041	16.83	0.0000		
					0.835	0.737
Volume						
Intercept ( $\alpha$ )	-0.35	0.077	-4.6	0.0000		
Slope ( $b$ )	0.91	0.052	17.3	0.0000		
					0.766	0.754

the full data set [i.e., the data include both the original and expanded (additional 18 observations) half-life data].

The prediction intervals in Figures 1 and 2 are supplemented by Tables 2 and 3, which present the intervals for a wide range of rat values in numerical form.

To illustrate the use and interpretation of the information contained in these tables, consider a newly encountered substance,  $X$ , whose human half-life we desire to ascertain. Say that from a toxicokinetic study in rats,  $X$  is estimated to exhibit a half-life of 0.8 hr. Table 2 provides the following information about the prediction of the human half-life. First, the prediction, or best guess of the human half-life is 4.02 hr. If, for example, we desired to have an interval in which we were 90% confident that the human half-life would lie, Table 3 indicates that 1.0–16.2 hr is the interval. That is, based on the results of this regression analysis, we can be 90% certain that the human half-life for  $X$  is between 1.0 and 16.2 hr. If interpolation is required, the table is constructed in such a way that linear interpolation should be adequate. For instance, if the rat half-life had been estimated to be 2.7 hr instead of 0.8, then interpolating linearly between the predicted human values of 8.60 (for a rat half-life of 2 hr) and 12.04 (for a rat half-life of 3 hr), the prediction for human half-life would be 11.01 hr.

## Discussion

In contrast to customary scaling studies, we wanted to determine whether a model could be developed for predicting the toxicokinetic or pharmacokinetic parameters of any xenobiotic in humans based expressly on the availability of estimates of those parameters in rats. To do so we regressed  $P_{\text{human}}$  against  $P_{\text{rat}}$  for over 100 xenobiotics. The data were accommodated by the model:

$$P_{i_{\text{human}}} = a P_{i_{\text{rat}}}^b \quad (5)$$

which, employing the logarithmic transformation, accounts for over 75% of the variance in the relationship between  $P_{\text{human}}$  and  $P_{\text{rat}}$ . This is particularly striking for a variety of reasons, including, though not limited to, the following: 1) multiple strains of rats were used in assimilating the data sets; 2) some of the data were acquired from male rats and some from female rats; 3) it was assumed, though not necessarily demonstrated, that elimination kinetics in all cases was first order. In fact, some of the xenobiotics exhibited clear dose-dependent kinetics in rats, and when such was the case the kinetic parameters were selected from the

**Table 2.** Table for predicting human half-life from rat half-life

Rat half-life	Lower			Human half-life estimate	Upper		
	95%	90%	80%		80%	90%	95%
0.01	0.019	0.025	0.034	0.106	0.327	0.451	0.598
0.02	0.034	0.045	0.062	0.189	0.574	0.790	1.045
0.03	0.048	0.064	0.087	0.264	0.799	1.098	1.450
0.04	0.062	0.081	0.111	0.335	1.011	1.387	1.830
0.05	0.074	0.098	0.134	0.404	1.213	1.664	2.193
0.06	0.087	0.114	0.156	0.469	1.408	1.930	2.543
0.07	0.099	0.130	0.178	0.533	1.598	2.189	2.882
0.08	0.110	0.146	0.199	0.596	1.782	2.441	3.213
0.09	0.122	0.161	0.220	0.657	1.963	2.687	3.536
0.1	0.13	0.18	0.24	0.72	2.14	2.93	3.85
0.2	0.24	0.31	0.43	1.27	3.78	5.17	6.79
0.3	0.34	0.44	0.60	1.78	5.28	7.21	9.47
0.4	0.43	0.56	0.77	2.26	6.70	9.14	12.00
0.5	0.51	0.68	0.92	2.72	8.05	10.99	14.42
0.6	0.60	0.79	1.07	3.17	9.36	12.77	16.76
0.7	0.68	0.89	1.22	3.60	10.63	14.51	19.04
0.8	0.76	1.00	1.36	4.02	11.88	16.20	21.26
0.9	0.84	1.10	1.50	4.44	13.09	17.86	23.44
1	0.92	1.20	1.64	4.84	14.29	19.49	25.57
2	1.63	2.13	2.91	8.60	25.40	34.66	45.47
3	2.27	2.98	4.07	12.04	35.59	48.58	63.76
4	2.88	3.78	5.16	15.28	45.24	61.76	81.09
5	3.46	4.54	6.20	18.39	54.49	74.42	97.74
6	4.02	5.28	7.21	21.39	63.46	86.69	113.88
7	4.56	5.99	8.19	24.31	72.18	98.64	129.61
8	5.08	6.68	9.14	27.15	80.71	110.32	144.99
9	5.60	7.36	10.06	29.94	89.07	121.77	160.08
10	6.1	8.0	11.0	32.7	97.3	133.0	174.9
20	10.7	14.1	19.4	58.1	174.0	238.3	313.9
30	14.9	19.7	27.0	81.3	244.6	335.6	442.4
40	18.8	24.9	34.1	103.2	311.6	427.9	564.7
50	22.6	29.8	41.0	124.1	376.1	516.9	682.7
60	26.2	34.6	47.5	144.4	438.6	603.3	797.2
70	29.6	39.2	53.9	164.1	499.5	687.5	909.1
80	33.0	43.6	60.1	183.3	559.2	770.0	1018.7
90	36.3	48.0	66.1	202.1	617.7	851.1	1126.4
100	39.5	52.3	72.0	220.6	675.2	930.8	1232.4
200	68.9	91.4	126.5	391.9	1214.2	1679.5	2230.5
300	95.2	126.7	175.7	548.6	1712.7	2374.3	3159.3
400	119.8	159.7	221.7	696.4	2186.8	3036.7	4046.6
500	143.2	191.0	265.6	837.9	2643.8	3676.1	4904.4
600	165.5	221.1	307.7	974.7	3087.5	4297.9	5739.6
700	187.1	250.1	348.5	1107.6	3520.5	4905.5	6556.7
800	208.1	278.3	388.1	1237.3	3944.6	5501.3	7358.6
900	228.5	305.8	426.8	1364.3	4361.2	6086.9	8147.4
1000	248.4	332.7	464.6	1488.9	4771.1	6663.7	8925.0
1100	267.9	359.0	501.7	1611.3	5175.1	7232.7	9692.5

lowest dose tested; 4) the sizes and ages of rats used were not uniform; 5) there was insufficient technical detail (e.g., sampling times, duration of sampling, number of subjects) to determine whether all studies from which kinetic data were gathered were structured appropriately (i.e., the goodness of the underlying data cannot be validated); 6) data for each xenobiotic were published generally without regard to chirality considerations; 7) data from a single experiment in a single sample of subjects were taken to be representative; 8) data for humans generally did not take account of sex; 9) no accommodation was made for the differences in protein-binding between species; 10) routes of administration were ignored;

11) elimination pathways were not identical for all xenobiotics in both species; and 12) when explicit weights were not provided, rat weights were set at 0.25 kg and human weights at 70 kg.

Inspection of Tables 2 and 3 shows that the 80% confidence limits for each predicted human parameter embrace an approximate 10-fold range of values. The full range of values for a given parameter for a given xenobiotic when estimated directly even from a single small sample of humans may also extend an order of magnitude or more. It is possible that the range of predicted values could have been tightened, and the scatter of data around the prediction line could have been diminished if



**Table 3.** Table for predicting human volume of distribution from rat volume of distribution

Rat volume	Lower			Human volume estimate	Upper		
	95%	90%	80%		80%	90%	95%
0.01	0.002	0.003	0.004	0.011	0.031	0.041	0.054
0.02	0.004	0.005	0.007	0.020	0.057	0.076	0.099
0.03	0.006	0.008	0.011	0.029	0.082	0.109	0.141
0.04	0.008	0.010	0.014	0.038	0.105	0.141	0.182
0.05	0.010	0.013	0.017	0.046	0.128	0.172	0.222
0.06	0.012	0.015	0.020	0.055	0.151	0.202	0.261
0.07	0.013	0.017	0.023	0.063	0.174	0.232	0.299
0.08	0.015	0.019	0.026	0.071	0.196	0.261	0.337
0.09	0.017	0.022	0.029	0.079	0.217	0.290	0.374
0.1	0.019	0.024	0.032	0.087	0.239	0.319	0.411
0.2	0.035	0.045	0.060	0.164	0.445	0.593	0.762
0.3	0.051	0.065	0.087	0.236	0.641	0.853	1.096
0.4	0.066	0.085	0.113	0.307	0.831	1.105	1.419
0.5	0.081	0.104	0.139	0.376	1.016	1.352	1.735
0.6	0.096	0.123	0.164	0.443	1.198	1.593	2.045
0.7	0.111	0.142	0.189	0.510	1.377	1.832	2.350
0.8	0.125	0.160	0.213	0.575	1.554	2.067	2.652
0.9	0.139	0.178	0.237	0.640	1.729	2.299	2.950
1	0.15	0.20	0.26	0.70	1.90	2.53	3.25
2	0.29	0.37	0.49	1.32	3.57	4.75	6.09
3	0.41	0.53	0.70	1.91	5.16	6.87	8.82
4	0.53	0.69	0.91	2.48	6.71	8.93	11.47
5	0.65	0.84	1.18	3.03	8.23	10.96	14.08
6	0.77	0.99	1.32	3.58	9.72	12.95	16.64
7	0.88	1.14	1.51	4.11	11.19	14.92	19.17
8	0.99	1.28	1.71	4.64	12.65	16.86	21.68
9	1.11	1.42	1.90	5.17	14.09	18.79	24.17
10	1.21	1.56	2.08	5.69	15.52	20.70	26.63
20	2.25	2.90	3.88	10.66	29.32	39.20	50.54
30	3.22	4.16	5.57	15.40	42.60	57.04	73.63
40	4.15	5.37	7.20	19.99	55.54	74.46	96.23
50	5.06	6.54	8.78	24.48	68.24	91.59	118.48
60	5.94	7.69	10.33	28.88	80.77	108.50	140.45
70	6.80	8.81	11.84	33.22	93.14	125.22	162.21
80	7.65	9.91	13.34	37.49	105.39	141.79	183.79
90	8.48	11.00	14.81	41.72	117.54	158.23	205.22

data sets had been developed prospectively rather than retrospectively and attention had been paid to the itemized points above.

The *sine qua non* for risk assessment purposes appears to be PBPK modeling, which mathematically describes uptake, distribution, metabolism, and excretion of

chemicals for physiologically relevant tissue compartments (11,12). However, 24 or more physiological, physicochemical, and enzyme kinetic parameters must be known for a given xenobiotic in a given animal model in order for PBPK models to accurately simulate the time course of xenobi-

otics in tissue compartments. Those active in the field point to the difficulty and expense associated with the development of comprehensive models (12).

The simple model presented here allows for the prediction of body burdens of xenobiotics during chronic exposure if the exposure rate is known:

$$\bar{A}_{ss} = E(1.44 \times t_{1/2}) \quad (6)$$

where  $\bar{A}_{ss}$  is the average amount of xenobiotic in the body at steady-state and  $E$  is the exposure rate (the fraction of dose absorbed during each exposure period). Alternatively, exposure rates can be estimated from measurement of an average steady-state plasma concentration:

$$E = FD/\tau = \bar{C}_{ss}(V/1.44) \times t_{1/2} \quad (7)$$

where  $D$  is the dose,  $F$  is the fraction of the dose absorbed,  $V$  is the volume of distribution,  $\tau$  is the interval of exposure, and  $\bar{C}_{ss}$  is the measured average steady-state concentration of the xenobiotic. To the extent that toxicity or carcinogenicity is a function of reactive metabolites, additional information would be required such as target tissue burdens of such metabolites at steady state. We do not propose that the simple model developed in this paper ought to supplant other strategies incorporated into risk assessment models, be used to predict initial doses of xenobiotics in humans, or be used to gauge the duration of exposures to xenobiotic challenges. Nevertheless, in an era of cost consciousness, it may be useful to apply simple strategies for such purposes when circumstances permit, and it may be instructive to ascertain those circumstances in which simple models may be as usefully informative as far more complex ones.

## Appendix

### Tabulation of Kinetic Parameters in Rats and Humans for 120 Xenobiotics

Xenobiotic	Volume of distribution (l/kg)				Half-life (hr)				Xenobiotic	Volume of distribution (l/kg)				Half-life (hr)			
	Rat	Ref.	Human	Ref.	Rat	Ref.	Human	Ref.		Rat	Ref.	Human	Ref.	Rat	Ref.	Human	Ref.
ALO 1567	0.45	(13)	0.77	(13)	15.59	(13)	63	(13)	Alteplase	0.08	(24)	0.05	(15)	0.04	(109)	0.07	(109)
ALO 1576	0.97	(13)	2.40	(13)	33.61	(13)	72	(13)	Amiodarone	72.3	(25)	66	(15)				
3TC <sup>a</sup>					1.6	(117)	2.6	(117)	Amitriptyline	15.83	(26)	15	(15)	1.47	(26)	15.06	(15)
Aspirin	0.37	(14)	0.15	(15)	0.07	(14)	0.29	(15)	Amobarbital	1.46	(27)	1.01	(27)	1.73	(27)	22.7	(27)
AZT <sup>a</sup>	2.12	(16)	1.40	(16)	0.39	(16)	1.35	(16)	Amoxicillin	0.42	(28)	0.21	(15)	0.3	(28)	1.7	(15)
CSF <sup>a</sup>					1.1	(118)	3.5	(119)	Amphotericin B	3.46	(29)	3.37	(15)	10.2	(29)	86	(125)
FCE 22101	0.05	(17)	0.24	(17)	0.09	(17)	0.74	(17)	Antipyrine	0.92	(30)	0.56	(31)	1.48	(28)	12	(110)
L-DOPA	9.8	(18)	1.65	(19)					Aprobarbital					5.79	(99)	24	(119)
PCP (phencyclidine)	23.2	(20)	6.20	(20)	2.2	(20)	16	(20)	Atenolol	3.41	(32)	0.95	(15)				
MK476 <sup>a</sup>					1.4	(120)	3.8	(120)	Baclofen	1.19	(33)	0.48	(34)				
Acetaminophen	1.02	(21)	0.95	(15)	0.27	(107)	2	(15)	Betamipron	0.24	(35)	0.28	(35)	0.16	(35)	0.81	(35)
Acivicin	0.65	(22)	0.50	(15)	1.44	(22)	9.5	(22)	HI-6 <sup>a</sup>					0.39	(121)	1.42	(121)
Acyclovir	0.7	(23)	0.69	(15)	1.39	(23)	2.4	(108)	Brodifacoum <sup>a</sup>					156	(123)	576	(122)
									Caffeine	0.92	(36)	0.61	(15)	0.85	(36)	4.9	(15)

appendix continued

## appendix, continued

Xenobiotic	Volume of distribution (l/kg)				Half-life (hr)				Xenobiotic	Volume of distribution (l/kg)				Half-life (hr)			
	Rat	Ref.	Human	Ref.	Rat	Ref.	Human	Ref.		Rat	Ref.	Human	Ref.	Rat	Ref.	Human	Ref.
Carbamazepine	12.91	(37)	1.4	(15)					Lithium <sup>a</sup>					6.1	(135)	28.9	(135)
Cefadroxil	0.79	(38)	0.22	(15)	0.82	(38)	1.1	(15)	Meberine <sup>a</sup>					0.48	(136)	2.5	(136)
Cefazolin	0.22	(39)	0.12	(15)	0.39	(39)	1.8	(15)	Methadone	7.81	(82)	4.24	(15)	1.45	(82)	35	(15)
Cefmetazole	0.49	(39)	0.14	(39)	0.14	(39)	1.55	(39)	Methotrexate	0.96	(51)	0.55	(15)	2.66	(51)	8.4	(51)
Cefodizime	0.18	(40)	0.19	(41)	1.82	(40)	1.6	(41)	Methylmercury <sup>a</sup>					1776	(137)	1056	(138)
Cefoperazone	0.29	(39)	0.09	(15)	0.15	(39)	2.1	(15)	Methysergide	1.43	(80)	0.93	(81)	0.02	(80)	0.75	(81)
Cefotetan	0.25	(39)	0.13	(15)	0.22	(39)	2.57	(15)	Metoclopramide	1.13	(83)	2.68	(15)	0.33	(83)	5	(15)
Cefpiramide	0.26	(39)	0.11	(39)	0.32	(39)	5.41	(39)	Metoprolol	6.72	(32)	4.16	(15)	0.66	(115)	3.2	(15)
Cefpirome	0.57	(42)	0.31	(43)					Midazolam	1.64	(84)	1.1	(15)	0.28	(84)	1.9	(15)
Ceftizoxime	0.44	(44)	0.36	(15)	0.26	(44)	1.8	(15)	Mitomycin C <sup>a</sup>					0.47	(139)	0.67	(139)
Cephadrine	0.43	(45)	0.25	(15)	0.81	(45)	0.77	(15)	Morphine	2.87	(85)	3.3	(15)	0.46	(116)	1.9	(15)
Chlorpromazine	29.1	(46)	11.2	(46)	5.55	(46)	30	(15)	Moxalactam	0.28	(39)	0.25	(15)	0.34	(39)	2.1	(15)
Cimetidine	1.36	(47)	1.0	(15)	0.4	(111)	2	(15)	Nicardipine	1.28	(86)	1.1	(15)	0.13	(86)	1.3	(15)
Cisplatin	2.06	(48)	0.28	(15)	0.79	(48)	0.53	(15)	Nicotine	5.70	(87)	2.6	(15)	1.16	(87)	2	(15)
Clavulanic acid	0.44	(49)	0.21	(15)					Ofloxacin	1.54	(88)	2.95	(89)				
Cocaine	17.2	(50)	2.0	(15)	0.85	(50)	0.8	(15)	Oxazepam <sup>a</sup>					4.5	(140)	6.8	(15)
Codeine	3.6	(50)	2.6	(15)	0.55	(124)	2.9	(15)	Panipenem	0.19	(35)	0.18	(35)				
Cyclophosphamide	0.9	(51)	0.78	(15)	2.49	(51)	5.25	(51)	Pentamidine <sup>a</sup>					0.03	(141)	0.28	(141)
Cyclosporine	4.54	(52)	1.2	(15)	15.51	(52)	16	(52)	Pentazocine	7.66	(90)	7.1	(15)	1.13	(90)	4.6	(15)
Diazepam	5.3	(53)	1.1	(15)	1.42	(53)	43	(15)	Pentobarbital	1.64	(27)	0.99	(27)	2.26	(27)	22.3	(27)
Dideoxycytidine					0.98	(126)	1.2	(127)	Phenobarbital	1.02	(27)	0.54	(15)	6.32	(27)	99	(15)
Diffunisal	0.13	(54)	0.10	(15)	1	(54)	11.55	(15)	Phenprocoumon <sup>a</sup>					17.4	(142)	128	(143)
Digoxin	3.48	(55)	9.19	(15)	2.5	(55)	39	(15)	Phenylbutazone	0.24	(27)	0.10	(15)	4.17	(27)	56	(15)
Enprofylline	0.34	(56)	0.51	(56)	0.27	(56)	1.42	(128)	Physostigmine					0.25	(144)	0.42	(145)
Erythropoietin <sup>a</sup>	0.07	(57)	0.03	(58)	2.64	(57)	5	(58)	Piroxicam	1.94	(91)	0.15	(15)	8.4	(91)	48	(91)
Ethanol	0.61	(59)	0.54	(15)					Panipenem					0.08	(35)	0.84	(35)
Ethosuximide	0.70	(60)	0.72	(15)	5.5	(129)	33	(130)	Prednisolone	1.74	(92)	1.5	(15)	0.2	(93)	2.2	(15)
Ethoxyacetic acid <sup>a</sup>					7.2	(131)	42	(131)	Prednisone	1.3	(93)	0.97	(15)	0.1	(93)	3.11	(15)
Etretinate	1.33	(61)	1.50	(62)					Procainamide	3.72	(94)	1.9	(15)	0.69	(94)	3	(15)
Famotidine	1.65	(63)	1.15	(15)	0.44	(63)	2.83	(63)	Propafenone	5.2	(95)	3.88	(95)	0.97	(95)	6.5	(95)
Felbamate	0.42	(64)	0.85	(65)	2.78	(64)	13.3	(64)	Propranolol	5.3	(96)	4.3	(15)	0.72	(96)	2.78	(96)
Fentanyl	11.2	(66)	7.7	(112)	1.93	(112)	7.9	(112)	Pyrimethamine	1.14	(97)	2.9	(15)				
Fluconazole <sup>a</sup>					4	(132)	26.4	(132)	Quinidine	6.0	(98)	2.7	(15)	2.05	(98)	6.2	(15)
Fluoxetine	15.9	(67)	35	(15)	3.92	(67)	42.1	(15)	Secobarbital	0.97	(99)	1.5	(100)	0.89	(99)	26.5	(100)
Flurbiprofen	0.36	(68)	0.21	(69)					Sertraline	23	(101)	25	(102)	4.5	(101)	24	(102)
Gabapentin <sup>a</sup>					2	(133)	5.5	(133)	Sulfadiazine	0.39	(103)	0.29	(15)	3.86	(103)	9.9	(15)
Gentamicin	0.6	(70)	0.31	(15)					Sulfizoxazole	0.32	(103)	0.15	(15)	4.44	(103)	6.6	(15)
Hexobarbital	0.7	(27)	1.2	(15)	0.42	(134)	4.4	(134)	Theophylline <sup>a</sup>	0.50	(79)	0.50	(15)	2	(146)	7.3	(146)
Ibuprofen	0.55	(71)	0.15	(15)	1.31	(71)	2	(15)	Ticarcillin	0.25	(49)	0.21	(15)	0.22	(49)	1.21	(15)
Imipramine	17.48	(72)	23	(72)	3.42	(113)	18	(15)	Tolbutamide	0.22	(27)	0.10	(15)	1.82	(27)	7.2	(27)
Indomethacin	0.12	(73)	0.26	(15)	2.77	(114)	2.4	(15)	Tolmetin	0.18	(104)	0.54	(15)				
Isoniazid	0.63	(74)	0.67	(15)	0.74	(74)	3.1	(15)	Trimethoprim	2.47	(105)	1.8	(15)	0.61	(105)	11	(15)
Isotretinoin	1.8	(75)	7	(15)					Valproate	0.66	(27)	0.13	(27)	4.6	(27)	14	(15)
Ketoconazole	0.66	(76)	2.4	(15)	0.53	(76)	3.3	(15)	Vancomycin	1.02	(106)	0.39	(15)	1.44	(106)	5.6	(15)
Ketoprofen	0.37	(77)	0.15	(15)					Verapamil					1.6	(147)	4	(15)
Ketorolac	0.38	(78)	0.11	(78)	1.8	(78)	6.5	(78)	Warfarin	0.20	(27)	0.14	(15)	11.1 <sup>b</sup>	(27,148)	37	(15)
Lidocaine	2.52	(79)	1.1	(15)	0.62	(79)	1.38	(15)									

<sup>a</sup>Denotes subset of xenobiotics tested against model that had been developed without the subset (see Results).<sup>b</sup>Unweighted average of mean values reported in Sawada et al. (27) and Takada and Levy (148).

## REFERENCES

- Boxenbaum H, Ronfeld R. Interspecies pharmacokinetic scaling and the Dedrick plots. *Am J Physiol* 245:R768–R775 (1983).
- Boxenbaum H. Interspecies pharmacokinetic scaling and the evolutionary-comparative paradigm. *Drug Metab Rev* 15:1071–1121 (1984).
- Gerlowski L, Jain R. Physiologically based pharmacokinetic modeling: principles and applications. *J Pharm Sci* 72:1103–1126 (1983).
- Sawada Y, Harashima H, Hanano M, Sugiyama Y, Iga T. Prediction of the plasma concentration time courses of various drugs in humans based on data from rats. *J Pharmacobiodyn* 8:757–766 (1985).
- Boxenbaum H. Interspecies scaling, allometry, physiological time, and the ground plan of pharmacokinetics. *J Pharmacokinet Biopharm* 10:201–227 (1982).
- Fan A, Howd R, Davis B. Risk assessment of environmental chemicals. *Annual Reviews of Pharmacol Toxicol* 35:341–368 (1995).
- Travis C, White R. Interspecies scaling of toxicity data. *Risk Anal* 8:119–125 (1988).
- Sidhu K. Basis for body weight exponent (0.75) as a scaling factor in energy metabolism and risk assessment. *J Appl Toxicol* 12:309–310 (1992).
- Travis C, White R, Ward R. Interspecies extrapolation of pharmacokinetics. *J Theor Biol* 142:285–304 (1990).
- Boxenbaum H, Fertig J. Scaling of antipyrine intrinsic clearance of unbound drug in 15 mammalian species. *Eur J Drug Metab Pharmacokinet* 9:177–183 (1984).
- Andersen ME, Krishnan K, Conolly R, McClellan R. Biologically based modeling in toxicology research. *Arch Toxicol* 15:217–227 (1992).
- Krewski D, Withey J, Ku L, Andersen ME. Applications of physiologic pharmacokinetic modeling in carcinogenic risk assessment. *Environ Health Perspect* 102(suppl 11):37–50 (1994).
- Brazzell R, Park T, Wooldridge C, McCue B, Barker R, Couch R, Yark B. Interspecies comparison of the pharmacokinetics of aldose reductase inhibitors. *Drug Metab Dispos* 18:435–440 (1990).
- Wientjes MG, Levy G. Nonlinear pharmacokinetics.

- netics of aspirin in rats. *J Pharmacol Exp Ther* 245:809–815 (1988).
15. Benet L, Williams R. Design and optimization of dosage regimens. In: *The pharmacological basis of therapeutics*, 8th ed (Goodman Gilman A, Rall T, Nies A, Taylor P, eds). New York: Pergamon Press, 1990;1650–1735.
  16. Patel B, Boudinot D, Schinazi R, Galb J, Chu C. Comparative pharmacokinetics and interspecies scaling of 3'-azido-3'-deoxythymidine (AZT) in several mammalian species. *J Pharmacobio-Dyn* 13:206–211 (1990).
  17. Efthymiopoulos C, Battaglia R, Strolin MB. Animal pharmacokinetics and interspecies scaling of FCE 22101, a penem antibiotic. *J Antimicrob Chemother* 27:517–526 (1991).
  18. Rose S, Jenner P, Marsder C. Peripheral pharmacokinetics handling and metabolism of L-dopa in the rat: the effect of route of administration and cardidopa pretreatment. *J Pharm Pharmacol* 43:325–330 (1991).
  19. Robertson DR, Wood ND, Everest H, Monks K, Waller DG, Renwick A, George CF. The effect of age on the pharmacokinetics of levodopa administered alone and in the presence of carbidopa. *Br J Clin Pharmacol* 28:61–69 (1989).
  20. Owens S, Hardwick W, Blackall D. Phencyclidine pharmacokinetic scaling among species. *J Pharmacol Exp Ther* 242:96–101 (1987).
  21. Belanger P, Lalande M, Dore F, Labricque G. Time-dependent variations in the organ extraction ratios of acetaminophen in rat. *J Pharmacokinet Biopharm* 15:133–141 (1987).
  22. McGovern J, Williams M, Stewart J. Interspecies comparison of acivicin pharmacokinetics. *Drug Metab Dispos* 16:18–22 (1988).
  23. Fujitaka Y, Mizuno N, Morita E, Motozono H, Takahashi K, Yamanaka Y, Shinkuma D. Effect of age on the gastrointestinal absorption of acyclovir in rats. *J Pharm Pharmacol* 43:465–469 (1991).
  24. Hori R, Ite Y, Shima T, Invi K, Aoki S, Okumura K, Tanigawa Y. Total body and hepatic clearance in rats of recombinant tissue type plasminogen activator expressed in mouse C127 and chinese hamster ovary cells. *Drug Metab Dispos* 20:541–546 (1992).
  25. Wyss P, Moor J, Bickel M. Single-dose kinetics of tissue distribution, excretion and metabolism of amiodarone in rats. *J Pharmacol Exp Ther* 254:502–504 (1990).
  26. Miyake K, Fukuchi H, Kitaura T, Kimura M, Kimura Y. Pharmacokinetic analysis of amitriptyline and demethylated metabolite in serum and brain of rats after acute and chronic oral administration of amitriptyline. *J Pharmacobio-Dyn* 15:157–166 (1992).
  27. Sawada Y, Hanano M, Sugiyama Y, Iga T. Prediction of the disposition of nine weakly acidic and six weakly basic drugs in humans from pharmacokinetics in rats. *J Pharmacokinet Biopharm* 13:477–492 (1985).
  28. Torres-Molina F, Aristorena J, Garcia-Carbonell, Granero L, Chese-Jimenez J, Pladelfina J, Peris-Ribeno J. Influence of permanent cannulation of the jugular vein on pharmacokinetics of amoxycillin and antipyrine in the rat. *Pharm Res* 9:1587–1591 (1992).
  29. Fielding R, Smith P, Wang L, Porter J, Guo L. Comparative pharmacokinetics of amphotericin B after administration of a novel coloidal delivery system, ABCB, and a conventional formulation to rats. *Antimicrob Agents Chemother* 35:1208–1213 (1991).
  30. Ben-Zvi Z, Hreash F, Kaplanski J. Disposition of hexobarbitone and antipyrine in DOCA-hypertensive rats. *J Pharm Pharmacol* 43:349–352 (1991).
  31. Shobha J, Raghuram T. Antipyrine kinetics in undernourished diabetics. *Eur J Clin Pharmacol* 41:359–361 (1991).
  32. Belpaire F, DeSmet F, Vyackier L, Vermeulen A, Rosseel M, Bogaert M, Chauvelot-Moachan L. Effect of aging on the pharmacokinetics of atenolol, metoprolol and propranolol in the rat. *J Pharmacol Exp Ther* 254:116–122 (1990).
  33. Mandema J, Heijligers-Feijen C, Tukker E, DeBoer A, Danhof M. Modeling of the effect site equilibration kinetics and pharmacodynamics of racemic baclofen and its enantiomers using quantitative EEG effect measures. *J Pharmacol Exp Ther* 261:88–94 (1992).
  34. Wuis E, Dirks M, Termond E, Vree T, Vanderkleijn E. Plasma and urinary excretion kinetics of oral baclofen in healthy subjects. *Eur J Clin Pharmacol* 37:181–184 (1989).
  35. Kurihara A, Naganuma H, Hisaoka M, Tokiwn H, Kawahara G. Prediction of human pharmacokinetics of panipenem-betamipron, a new carbapenem from animal data. *Antimicrob Agents Chemother* 36:1810–1815 (1992).
  36. Bonati M, Latini R, Tognoni G, Young J, Garattini S. Interspecies comparison of in vivo caffeine pharmacokinetics in man, monkey, rabbit, rat and mouse. *Drug Metab Rev* 15:1355–1383 (1984–85).
  37. Turner P, Renton K. The interaction between carbamazepine and erythromycin. *Can J Physiol Pharmacol* 67:582–586 (1989).
  38. Garcia-Carborell M, Granero L, Torres-Molina F, Aristorena J, Peris-Ribera E. Nonlinear pharmacokinetics of cefadroxil in the rat. *Drug Metab Dispos* 21:215–217 (1993).
  39. Sawada Y, Hanano M, Sugiyama Y, Iga T. Prediction of the disposition of  $\beta$ -lactam antibiotics in humans from pharmacokinetic parameters in animals. *J Pharmacokinet Biopharm* 12:241–261 (1984).
  40. Matsushita H, Suzuki H, Sugiyama Y, Sawada Y, Iga T, Hanano M, Kawaguchi Y. Prediction of the pharmacokinetics of cefodizime and cefotetan in humans from their pharmacokinetic parameters in animals. *J Pharmacobio-Dyn* 13:602–611 (1990).
  41. Korting H, Schafer-Korting M, Maass L, Klesel N, Mutschler E. Cefodizime in serum and skin blister fluid after single intravenous and intramuscular doses in healthy volunteers. *Antimicrob Agents Chemother* 31:1822–1825 (1987).
  42. Isert D, Klesel N, Limbert M, Markus A, Seibert G, Schrinner E. Pharmacokinetics of cefiprone administered intravenously or intramuscularly to rats and dogs. *J Antimicrob Chemother* 29:31–37 (1992).
  43. Kavi J, Andrews J, Ashby J, Hillman G, Wise R. Pharmacokinetics and tissue penetration of ceftiprome, a new cephalosporin. *J Antimicrob Chemother* 22:911–916 (1988).
  44. Mordenti J. Pharmacokinetic scale-up: accurate prediction of human pharmacokinetic profiles from animal data. *J Pharm Sci* 74:1098 (1995).
  45. Nakashima E, Matshushita R, Takeaki M, Nakanishi T, Ichimura F. Comparative pharmacokinetics of cefoperazone and cephadrine in untreated streptozotocin diabetic rats. *Drug Metab Dispos* 20:730–735 (1992).
  46. Pettijohn F, Potter P, Siever R. Relation between binding to plasma protein, apparent volume of distribution and rate constants of disposition and elimination for chlorpromazine in three species. *J Pharm Pharmacol* 24:818–819 (1972).
  47. Gomita Y, Eto K, Furuno K, Araki Y. Effects of exposure to standard-and-nicotine-reduced cigarette smoke on pharmacokinetics of theophylline and cimetidine in rats. *J Pharm Sci* 81:1132–1135 (1992).
  48. Litterst C, Magin R. Alterations in plasma pharmacokinetics of cisplatin in tumor-bearing rats. *Cancer Chemother Pharmacol* 22:1–4 (1988).
  49. Woodnutt G, Kernutt I, Mizen L, Mizen L. Pharmacokinetics and distribution of ticarcillin-clavulanic acid (Timentin) in experimental animals. *Antimicrob Agents Chemother* 31:1826–1830 (1987).
  50. Boni J, Barr W, Martin B. Cocaine inhalation in the rat: pharmacokinetics and cardiovascular response. *J Pharmacol Exp Ther* 257:307–315 (1991).
  51. Boxenbaum H. Interspecies scaling, allometry, physiological time and the ground plan of pharmacokinetics. *J Pharmacokin Biopharm* 10:206–213 (1982).
  52. Sangalli L, Bartolotti, Jiritano L, Bonati M. Cyclosporine pharmacokinetics in rats and interspecies comparison in dogs, rabbits, rats and humans. *Drug Metab Dispos* 16:749–753 (1988).
  53. Igari Y, Sugiyama Y, Sawada Y, Iga T, Hanano M. In vitro and in vivo assessment of hepatic and extra-hepatic metabolism of diazepam in the rat. *J Pharm Sci* 73:825–828 (1984).
  54. Lin J, Hooke K, Yeh K, Duggan D. Dose-dependent pharmacokinetics of diflunisal in rats: dual effects of protein binding and metabolism. *J Pharmacol Exp Ther* 235:402–406 (1985).
  55. Harrison L, Gibaldi M. Pharmacokinetics of digoxin in the rat. *Drug Metab Dispos* 4:88–93 (1976).
  56. Tsunekawa Y, Hasegawa T, Nadai M, Takagi K, Nabeshima T. Interspecies differences and scaling for the pharmacokinetics of xanthine derivatives. *J Pharm Pharmacol* 44:594–599 (1992).
  57. Kinoshita H, Ohishi N, Kato M, Tokura S, Okazaki A. Pharmacokinetics and distribution of recombinant erythropoietin in rats. *Arzneim-Forsch* 42:174–178 (1992).
  58. Salmonson T, Danielson B, Wikstrom B. The pharmacokinetics of recombinant human erythropoietin after intravenous and subcutaneous administration to healthy subjects. *Br J Clin Pharmacol* 29:709–713 (1990).
  59. DaSilva V, McLean A. Effect of two different types of malnutrition on the rate of elimination of ethanol in rats. *Biochem Pharmacol* 37:4235–4238 (1988).
  60. Bachmann K, Jahn D, Yang C, Schwartz J. Ethosuximide disposition kinetics in rats. *Xenobiotica* 18:373–380 (1988).
  61. Chien D, Sandri R, Tang-Liu D. Systemic pharmacokinetics of acitretin, etretinate, isotretinoin and acetylenic retinoids in guinea

- pigs and obese rats. *Drug Metab Dispos* 20:211–217 (1992).
62. Larsen F, Jakobsen P, Larsen C, Norgaard A, Kragballe K, Nielsen-Kudsk F. Single dose pharmacokinetics of etretin and etretinate in psoriatic patients. *Pharmacol Toxicol* 61:85–88 (1987).
  63. Lin JH, Los L, Ulm E, Duggan P. Urinary excretion kinetics of famotidine in rats. *Drug Metab Dispos* 15:212–215 (1987).
  64. Adusumalli V, Yang J, Wong K, Kucharczyk N, Sofia R. Felbamate Pharmacokinetics in the rat, rabbit and dog. *Drug Metab Dispos* 19:1116–1125 (1991).
  65. Graves N, Ludden T, Holmes G, et al. Pharmacokinetics of felbamate, a novel antiepileptic drug: Application of mixed-effect modeling to clinical trials. *Pharmacotherapy* 9:372–376 (1989).
  66. Hug C Jr, Murphy M. Tissue redistribution of fentanyl and termination of its effects in rats. *Anesthesiology* 55:369–375 (1981).
  67. Caccia S, Cappi M, Fracasso C, Garattini S. Influence of dose and route of administration on the kinetics of fluoxetine and its metabolite norfluoxetine in the rat. *Psychopharmacology* 100:509–514 (1990).
  68. Berry B, Jamali F. Enantiomeric interaction of flurbiprofen in the rat. *J Pharm Sci* 78:632–634 (1989).
  69. Kean W, Antal E, Grace E, Cauvier H, Rischke J, Buchanan W. The pharmacokinetics of flurbiprofen in younger and elderly patients with rheumatoid arthritis. *J Clin Pharmacol* 32:41–48 (1992).
  70. Engineer M, Bodley G Sr, Newman R, W-Ho D. Effects of cisplatin-induced nephrotoxicity on gentamicin pharmacokinetics in rats. *Drug Metab Dispos* 15:329–334 (1987).
  71. Shah A, Jung D. Dose-dependent pharmacokinetics of ibuprofen in the rat. *Drug Metab Dispos* 15:151–154 (1987).
  72. Sawada Y, Hanano M, Sugiyama Y, Harashima H, Iga T. Prediction of the volumes of distribution of basic drugs in human based on data from animals. *J Pharmacokin Biopharm* 12:587–596 (1984).
  73. Ogiso T, Ito Y, Iwaki M, Atago H. A pharmacokinetic model for the percutaneous absorption of indomethacin and the prediction of drug disposition kinetics. *J Pharm Sci* 78:319–323 (1989).
  74. Zeruesenay D, Siegmund W, Zschesche M, Franke G, Walter R, Bleyer H. Pharmacokinetic interactions between isoniazid and theophylline in rats. *J Pharm Pharmacol* 44:893–897 (1992).
  75. Chien D, Sandri R, Tang-Liu D. Systemic pharmacokinetics of acitretin, etretinate, isotretinoin and acetylenic retinoids in guinea pigs and obese rats. *Drug Metab Dispos* 20:211–217 (1992).
  76. Rimmel R, Amoh K, Abdel-Monem M. The disposition and pharmacokinetics of ketoconazole in the rat. *Drug Metab Dispos* 15:735–739 (1987).
  77. Foster R, Jamali F. Stereoselective pharmacokinetics of ketoprofen in the rat. *Drug Metab Dispos* 16:623–626 (1988).
  78. Mrosczak E, Lee F, Combs D, Sarquist F, Huang B, Wu A, Tokes L, Maddox M, Cho D. Ketorolac, tromethamine absorption, distribution, metabolism, excretion and pharmacokinetics in animals and humans. *Drug Metab Dispos* 15:618–626 (1987).
  79. Fruncillo R, Digregorio G. Pharmacokinetics of pentobarbital, quinidine, lidocaine and theophylline in the thermally injured rat. *J Pharm Sci* 73:1117–1121 (1984).
  80. Bredberg U, Paalzow L. Pharmacokinetics of methysergide and its metabolite methylergometrine in the rat. *Drug Metab Dispos* 18:338–343 (1990).
  81. Bredberg U, Eyjolfssdottir, Paalow L, Tfelt-Hansen V, Tfelt-Hansen V. Pharmacokinetics of methysergide and its metabolite methylergometrine in man. *Eur J Clin Pharmacol* 30:75–77 (1986).
  82. Ling G, Umans J, Inturrisio C. Radio immunoassay and pharmacokinetics in the rat. *J Pharmacol Exp Ther* 217:147–151 (1981).
  83. Bakke O, Segura J. The absorption and elimination of metoclopramide in three animal species. *J Pharm Pharmacol* 28:32–39 (1976).
  84. Aarons L, Mandema J, Danhof M. A population analysis of the pharmacokinetics and pharmacodynamics of midazolam in the rat. *J of Pharmacokin Biopharm* 19:485–496 (1991).
  85. Bhargava H, Villar V. Pharmacodynamics and pharmacokinetics of intravenously administered morphine in spontaneously hypertensive and normotensive Wistar-Kyoto rats. *J Pharmacol Exp Ther* 261:290–296 (1992).
  86. Higuchi S, Shiobana Y. Comparative pharmacokinetics of nicardipine hydrochloride, a new vasodilator in various species. *Xenobiotica* 10:447–454 (1980).
  87. Kveremater G, Taylor L, DeBethizv J, Vessel E. Pharmacokinetics of nicotine and 12 metabolites in the rat. Application of a new radiometric high performance liquid chromatographic assay. *Drug Metab Dispos* 16:125–129 (1988).
  88. Okezaki E, Terasaki T, Nakamura M, Nagata O, Kato H, Tsuji A. Structure-tissue distribution relationship based on physiological pharmacokinetics for NY-198, a new antimicrobial agent, and the related pyridonecarboxylic acids. *Drug Metab Dispos* 16:865–874 (1988).
  89. Lode H, Hoffken G, Prinzing C, Glatzel P, Wiley R, Olschewski P, Sievers B, Reimnitz D, Borner K, Koeppe P. Comparative pharmacokinetics of new quinolones. *Drugs* 34(suppl 1):21–25 (1987).
  90. Ichimura F, Yokoyawa K, Yamana T, Tsuji A, Mizukami Y. Physiologic pharmacokinetic model for pentazocin. I. tissue distribution and elimination in the rat. *Int J Pharm* 15:321–333 (1983).
  91. Said S, Foda A. Influence of cimetidine on the pharmacokinetics of piroxicam in rat and man. *Arzneim-Forsch* 39:790–792 (1989).
  92. Nicols A, D'Ambrosio R, Pysczynski N, Jusko W. Pharmacokinetics and pharmacodynamics of prednisolone in obese rats. *J Pharmacol Exp Ther* 250:963–970 (1989).
  93. Huang M, Jusko W. Nonlinear pharmacokinetics and interconversion of prednisolone and prednisone in rats. *J Pharmacokin Biopharm* 18:401–421 (1990).
  94. Basseches P, DiGregorio G. Pharmacokinetics of procainamide in rats with extra-hepatic biliary obstruction. *J Pharm Sci* 71:1256–1259 (1982).
  95. Puigdemont A, Guitart R, DeMora F, Arboix M. Prediction of the disposition of propafenone in humans and dogs from pharmacokinetic parameters in other animal species. *J Pharm Sci* 80:1106–1109 (1991).
  96. Evans G, Nies A, Shand D. The disposition of propranolol III. Decreased half-life and volume of distribution as a result of plasma binding in man, monkey, dog and rat. *J Pharmacol Exp Ther* 186:114–122 (1973).
  97. Mihaly G, Date N, VeeNendaal J, Newman K, Smallwood R. Decreased hepatic elimination of pyrimethamine during malaria infection. *Biochemical Pharmacol* 36:2827–2829 (1987).
  98. Harashima H, Sawada, Sugiyama Y, Iga T, Hanano M. Analysis of non-linear tissue distribution of quinidine in rats by physiologically based pharmacokinetics. *J Pharmacokin Biopharm* 13:425–440 (1985).
  99. Yih T, Van Rossum, J. Pharmacokinetics of some homologous series of barbiturates in the intact rat and in the isolated perfused rat liver. *J Pharmacol Exp Ther* 203:184–202 (1977).
  100. Breimer D. Clinical pharmacokinetics of hypnotics. *Clin Pharmacokin* 2:93–109 (1977).
  101. Tremaine L, Welch W, Ronfeld R. Metabolism and disposition of the 5-hydroxytryptamine uptake blocker sertraline in the rat and dog. *Drug Metab Dispos* 17:542–550 (1989).
  102. DeVane C. Pharmacokinetics of the selective serotonin reuptake inhibitors. *J Clin Psychiatry* 53:2(suppl) 13–19 (1992).
  103. Kaul S, Ritschel W. Quantitative structure-pharmacokinetic relationship of a series of sulfonamides in the rat. *Eur J Drug Metab Pharmacokin* 15:211–217 (1990).
  104. Sabater J, Domenech J, Obach R. Pharmacokinetic study of tolmetin in the rat. *Arzneim-Forsch* 42:950–953 (1992).
  105. Tu Y, Allen L, Fiorica V, Albers D. Pharmacokinetics of trimethoprim in the rat. *J Pharm Sci* 78:556–560 (1989).
  106. Engineer M, Ho D, Bodey G. Comparison of vancomycin disposition in rats with normal and abnormal renal function. *Antimicrob Agents Chemother* 20:718–722 (1981).
  107. Galinsky R, Levy G. Dose and time-dependent elimination of acetaminophen in rats: pharmacokinetic implications of cosubstrate depletion. *J Pharmacol Exp Ther* 219:14–20 (1981).
  108. Brigden D, Whiteman P. The clinical pharmacology of acyclovir and its prodrugs. *Scand J Infect Dis* 47(suppl)33–39 (1985).
  109. Tanswell P, Heinzl G, Greischel A, Krause J. Nonlinear pharmacokinetics of tissue-type plasminogen activator in three animal species and isolated perfused rat liver. *J Pharmacol Exp Ther* 255:318–324 (1990).
  110. Roberts R, Grice J, Wood L, Petroff V, McGuffie C. Cimetidine impairs the elimination of theophylline and antipyrine. *Gastroenterology* 81:19–21 (1981).
  111. Adedoyin A, Aarons L, Houston J. Dose-dependent pharmacokinetics of cimetidine in the rat. *Xenobiotica* 17:595–604 (1987).
  112. Bjorkman S, Stanski D, Verotta D, Harashima H. Comparative tissue concentration profiles of fentanyl and alfentanil in humans predicted from tissue/blood partition data obtained in rats. *Anesthesiology* 72:865–873 (1990).
  113. Taylor I, Ioannides C, Sacra P, Turner J, Parke D. Effect of deuteration of imipramine on its pharmacokinetic properties in the rat. *Biochem Pharmacol* 32:641–647 (1983).
  114. Guissou P, Cuisinaud G, Legheand J, Sassard J. Chronopharmacokinetics of indomethacin in rats. *Arzneim-Forsch* 37:1034–1037 (1987).

115. Vermeulen A, Belpaire F, DeSmet F, Vercruysse I, Bogaert M. Aging and the pharmacokinetics and metabolism of metoprolol enantiomers in the rat. *J Gerontol Biol Sci* 48:B108–B114 (1993).
116. Miyamoto Y, Ozaki M, Yamamoto H. Effects of adrenalectomy on pharmacokinetics and antinociceptive activity of morphine in rats. *Jpn J Pharmacol* 46:379–386 (1988).
117. Hussey E, Donn K, Daniel M, Hall S, Harker A, Evans G. Interspecies scaling and pharmacokinetic parameters of 3TC in humans. *J Clin Pharmacol* 34:975–977 (1994).
118. Tanaka H, Kaneko T. Pharmacokinetics of recombinant human granulocyte colony-stimulating factor in the rat. Single and multiple dosing studies. *Drug Metab Dispos* 19:200–204 (1991).
119. Drug facts and comparisons. St. Louis, MO: Facts and Comparisons, Inc., 1995.
120. Merck and Company. Clinical investigator's confidential informational brochure. MK-0476 summary of information. Rahway, NJ, 1994.
121. Baggot J. Application of interspecies scaling to the bispyridinium oxime HI-6. *Am J Vet Res* 55:689–691 (1994).
122. Hollinger B, Pastoor T. Case management and plasma half-life in a case of brodifacoum poisoning. *Arch Intern Med* 153:1925–1928 (1993).
123. Bachmann K, Sullivan T. Disposition and Pharmacodynamic characteristics of brodifacoum in warfarin-sensitive rats. *Pharmacology* 27:281–288 (1983).
124. Shah J, Mason W. A dose-ranging study of the pharmacokinetics of codeine phosphate following intravenous administration to rats. *J Pharm Sci* 80:229–231 (1991).
125. Sanders S, Buchi K, Goddard M, Lang J, Tolman K. Single-dose pharmacokinetics and tolerance of a cholesteryl sulfate complex of amphotericin B administered to healthy volunteers. *Antimicrob Agents Chemother* 35:1029–1034 (1991).
126. Ibrahim S, Boudinot F. Pharmacokinetics and 2', 3'-dideoxycytidine after high-dose administration to rats. *J Pharm Sci* 80:36–38 (1991).
127. Ibrahim S, Boudinot F. Pharmacokinetics and 2', 3'-dideoxycytidine in rats: application to interspecies scale-up. *J Pharm Pharmacol* 41:829–834 (1988).
128. Borga O, Andersson K, Edholm L, Fagerstrom P, Lunell E, Persson C. Enprofylline genetics in healthy subjects after single doses. *Clin Pharmacol Ther* 34:799–804 (1983).
129. Bachmann K, Chu C, Greear V. In vivo evidence that ethosuximide is a cytochrome P450IIIa substrate. *Pharmacology* 45:121–128 (1992).
130. Bachmann K, Schwartz J, Sullivan T, Jauregui L. Single sample estimate of ethosuximide clearance. *Int J Clin Pharmacol Ther Toxicol* 24:546–550 (1986).
131. Groeseneken D, Veulemans H, Masschelein R, Van Vlem E. Comparative urinary excretion of ethoxyacetic acid in man and rat after single low doses of ethylene glycol monoethyl ether. *Toxicol Lett* 41:57–68 (1988).
132. Jezequel S. Fluconazole: Interspecies scaling and allometric relationships of pharmacokinetic properties. *J Pharm Pharmacol* 46:196–199 (1993).
133. Vollmer K, von Hodenberg A, Kölle E. Pharmacokinetics and metabolism of gabapentin in rat, dog and man. *Arzneim-Forsch* 36:830–838 (1986).
134. Van der Graaff M, Vermeulen N, Breimer D. Disposition of hexobarbital: 15 years of an intriguing model substrate. *Drug Metab Rev* 19:109–164 (1988).
135. Wood A, Goodwin G, DeSouza R, Green A. The pharmacokinetic profile of lithium in rat and mouse. An important factor in psychopharmacological investigation of the drug. *Neuropharmacol* 25:1285–1288 (1986).
136. Dickinson R, Baker P, Franklin M, Hooper W. Facile hydrolysis of meberine in vitro and in vivo: Negligible circulating concentrations of the drug after oral administration. *J Pharm Sci* 80:952–957 (1991).
137. Farris F, Dedrick R, Allen P, Smith C. Physiological model for the pharmacokinetics of methylmercury in the growing rat. *Toxicol Appl Pharmacol* 119:74–90 (1993).
138. Smith J, Allen P, Turner M, Most B, Fisher H, Hall L. The kinetics of intravenously administered methylmercury in man. *Toxicol Appl Pharmacol* 128:251–256 (1994).
139. Malviya V, Young J, Boike G, Gove N, Deppe G. Pharmacokinetics of mitomycin-C in plasma and tumor tissue of cervical cancer patients and in selected tissues of female rats. *Gynecol Oncol* 25:160–170 (1986).
140. Yuan J, Goehl T, Hong L, Clark J, Murrill E, Moore R. Toxicokinetics of oxazepam in rats and mice. *J Pharm Sci* 83:1373–1379 (1994).
141. Bernard E, Donnelly H, Maher M, Armstrong D. Use of a new bioassay to study pentamidine pharmacokinetics. *J Infect Dis* 152:750–754 (1985).
142. Trenk D, Jähnchen E. Effect of serum protein binding on pharmacokinetics and anticoagulant activity of phenprocoumon in rats. *J Pharmacokinect Biopharm* 8:177–191 (1980).
143. Hausteink K, Huller G. Pharmacokinetics of phenprocoumon. *Int J Clin Pharmacol Ther* 32:192–197 (1994).
144. Somani S, Khalique A. Pharmacokinetics and pharmacodynamics of physostigmine in the rat after intravenous administration. *Drug Metab Dispos* 15:627–633 (1987).
145. Johansson M, Nordberg A. Pharmacokinetic studies of cholinesterase inhibitors. *Acta Neurol Scand* 149:522–525 (1993).
146. Gascón A, Calvo B, Hernandez R, Dominguez-Gil A, Pedraz J. Interspecies scaling of cimetidine-theophylline pharmacokinetic interaction: interspecies scaling in pharmacokinetic interactions. *Pharm Res* 11:945–950 (1994).
147. Manipitsitkul P, Chiou W. Intravenous verapamil kinetics in rats. Marked arteriovenous concentration difference and comparison with humans. *Biopharm Drug Dispos* 14:555–566 (1993).
148. Takada K, Levy G. Comparative pharmacokinetics of coumarin anticoagulants XLIV. Dose dependent pharmacokinetics of warfarin in rats. *J Pharm Sci* 69:9–14 (1980).

## CAAT Recognition Award

The Johns Hopkins Center for Alternatives to Animal Testing (CAAT) would like to honor an individual or organization who has made an outstanding contribution to the field of 3Rs alternatives and in vitro sciences. We invite the readers of this journal to submit nominations. The award will be presented at the second World Congress on Alternatives and Animal Use in the Life Sciences, to be held in October 1996 in Utrecht, The Netherlands. Deadline for receipt of nominations is June 1, 1996. Please send your nomination, including a one-page description of why this individual or organization should be recognized. Please include a curriculum vitae for individual nominees and a fact sheet or supporting documents for organizations. A subcommittee of the CAAT Advisory board will review the nominations and select the recipient of the CAAT Recognition Award.

Forward nominations to: Alan M. Goldberg, Ph.D., Johns Hopkins Center for Alternatives to Animal Testing  
111 Market Place, Suite 840, Baltimore, MD 21202-6709